



Applied StemCell, Inc.

(866) 497-4180

www.appliedstemcell.com

Datasheet

Substrate-free Autobioluminescent Cell Lines Human Colorectal Cancer Cells (HCT116)

Product Information

Catalog Number	ASE-5901
Description	Applied StemCell, Inc.'s Autobioluminescent HCT116 cell line is a substrate-independent synthetic luciferase system that encodes all of the components required for bioluminescent signal initiation and maintenance. This tumor forming line can be grown either two dimensionally or three dimensionally to fit a wide variety of assays and experimental designs, including <i>in vivo</i> applications requiring monitoring of tumor growth in small animal models. This is an ideal cell line for preclinical drug screening and discovery.
Quantity	1 x 10 ⁶ cells/ 1 mL
Cellular Morphology	Epithelial
Gender	Male
Tissue	Colon
Disease	Colorectal Carcinoma
Growth Properties	Adherent

Recommended Growth Medium

The following medium is recommended for cellular growth (not provided with cells)

Component	Vendor	Cat. Number	Final Conc.
HyClone™ McCoy's 5A medium	GE Life Sciences	SH30270	
ES-Sure™ FBS	Applied StemCell	ASM-5017	10%
100 mM Sodium Pyruvate	ThermoFisher	11360070	1 mM
HyClone™ Penicillin-Streptomycin 100X solution	GE Lifesciences	SV30010	1X
G 418 Sulfate, Sterile-Filtered Aqueous Solution	EMD Millipore	345812	200 µg/mL

Protocol

1. Thawing the Frozen Cells

- Incubate the vial in a 37°C water bath with gentle agitation until the contents of the vial have thawed (thawing should occur in approximately 2 minutes). To reduce the possibility of contamination, do not submerge the O-ring or cap during the thawing process.
- Spray the thawed vial with 70% ethanol and transfer to a sterilized environment. All remaining steps should be performed using aseptic techniques.

Applied StemCell, Inc.

521 Cottonwood Dr. #111, Milpitas, CA 95035

Phone: 866-497-4180 (US Toll Free); 408-773-8007 Fax: 408-773-8238

info@appliedstemcell.com www.appliedstemcell.com

Copyright 2016, Applied StemCell, Inc. All rights reserved. This information is subject to change without notice.

- 1.3. Transfer the full contents of the cryovial to a 15 mL conical tube containing 9 mL warmed (37°C) complete growth medium, and centrifuge at 125 x g for 7 minutes.
- 1.4. Resuspend the pellet in complete growth medium (warmed to 37°C) and transfer to an appropriate cell culture container.
Note: 25 cm² or 75 cm² flasks are recommended for initial thawing procedures.
- 1.5. Incubate the cells at 37°C and 5% CO₂ in a humidity controlled environment and monitor for growth.

2. Routine Growth and Maintenance

- 2.1. Remove and discard spent culture medium.
- 2.2. Rinse cells with an appropriate volume of sterile, 37°C PBS.
- 2.3. Add an appropriate volume of Trypsin-EDTA solution to the flask and incubate until cells have detached (detachment usually occurs within 2 to 15 minutes depending on if the incubation is performed at room temperature or at 37°C).
- 2.4. Resuspend the detached cells in an appropriate amount of pre-incubated 37°C complete growth medium.
- 2.5. Aliquot cells into new culture vessels containing pre-incubated 37°C complete growth medium.
- 2.6. Incubate the cells at 37°C and 5% CO₂ in a humidity controlled environment.

Subcultivation ratio: It is recommended that cells be subcultured at a ratio between 1:3 and 1:10 as needed.

Medium Refreshment: It is recommended that medium be refreshed every 2-3 days as needed.

Precautions and Disclaimers

Safety Precaution

PLEASE READ BEFORE HANDLING ANY FROZEN VIALS. These cells should be treated as biosafety level 1. Please wear the appropriate Personal Protective Equipment (lab coat, thermal gloves, safety goggles and a face shield) when handling the cells. Handle the frozen vials with due caution. Please be aware that the following scenario may occur: Liquid nitrogen may leak into the vials when the vials are submerged in liquid nitrogen. Upon thawing, the liquid nitrogen returns to the gas phase, resulting in a dangerous build-up of pressure within the vial. This can result in the vial exploding and expelling not only the vial contents but also the vial cap and plastic fragments of the vial.

Limited Use Label License

This product is to be used for internal, non-commercial research purposes for the sole benefit of the purchaser. It may not be used for any other purpose, including, but not limited to diagnostics or therapeutics, and may not be used in humans. This product may not be transferred or sold to third parties, resold, modified for resale, or used to manufacture or develop commercial products or to provide a service of any kind to third parties, including, without limitation, reporting the results of purchaser's activities for a fee or other form of consideration. For information on obtaining commercial research or additional rights, please contact Applied StemCell, Inc.

Disclaimer

Applied StemCell, Inc. makes every effort to include accurate and up-to-date information on this product information sheet. However, Applied StemCell makes no warranties or representation of its accuracy. This product is sent with the condition that you are responsible for its safe storage, handling, and use. Applied StemCell is not liable for any damages or injuries arising from the receipt and/or use of this product.