



## Neurons Starter Kit (iPSC from Blood Cells; Female)

### Product Information

<b>Catalog Number</b>	<b>ASE-9321KF (Female)</b>
<b>Description</b>	Applied StemCell's Neurons Starter Kit contains cryo-preserved, pre-differentiated mixed population neurons derived from a footprint-free, karyotype normal human iPSC line. It is designed for customers to generate mature neurons using Applied StemCell's optimized Neuron Maturation Medium and Supplements. Mature neurons can be obtained within 8 days. The neurons can be seeded on various culture vessel formats including 96-well plates on either glass or plastic surfaces and cultured as adherent cells. Shortly after seeding, the cells proliferate slightly for up to 3 days and show extensive neurite outgrowth and proper neuronal morphology. In general, on Day 8 post-seeding, the cell population will contain >90% neurons and <5% Glial Fibrillary Acidic Protein (GFAP) positive astrocytes.
<b>Characterization of Mature Neurons</b>	Maturation of neurons can be assessed by their morphology and by immunostaining of the Tuj-1 marker (neuronal class III- $\beta$ tubulin). Percentage of neurons can be determined by a count of Tuj-1 positive neurons divided by the total number of cells (DAPI staining of nuclei).
<b>Shipping</b>	Dry ice and at ambient temperature
<b>Safety Precaution</b>	<b>PLEASE READ BEFORE HANDLING ANY FROZEN VIALS.</b> Please wear the appropriate Personal Protection Equipment (lab coat, thermal gloves, safety goggles and a face shield) when handling the cells. Handle the frozen vials with due caution. Please be aware that the following scenario can occur: Liquid nitrogen can leak into the vials when the vials are submerged in liquid nitrogen. Upon thawing, the liquid nitrogen returns to the gas phase, resulting in a dangerous build-up of pressure within the vial. This can result in the vial exploding and expelling not only the vial contents but also the vial cap and plastic fragments of the vial.
<b>Restricted Use</b>	This product is for research use only and not intended for human or animal diagnostic or therapeutic uses.
<b>Warranty</b>	Performance of Applied StemCell's neurons has been extensively tested with other components of the Neurons Starter Kit. Applied StemCell will not hold responsibility if components other than those provided with Neurons Starter Kit are used to culture astrocyte precursors.

#### Applied StemCell, Inc.

521 Cottonwood Dr. #111, Milpitas, CA 95035  
Phone: 866-497-4180 (US Toll Free); 408-773-8007 Fax: 408-773-8238  
[info@appliedstemcell.com](mailto:info@appliedstemcell.com) [www.appliedstemcell.com](http://www.appliedstemcell.com)

**Media and Material**

**Neurons Starter Kit (ASE-9321KF)**

Component	Amount	Storage	Shelf Life
Neurons	1 vial $\geq 1 \times 10^6$ viable cells	Liquid Nitrogen	12 months
Neuron Maturation Medium	1 x 50 mL	-20°C	12 months
Supplement A	1 x 50 $\mu$ L vial	-20°C	12 months

**Additional Reagents Required**


- Recommended:
  - Poly-L-ornithine hydrobromide, Sigma, Cat# P3655
- Primary antibodies:
  - Mouse anti- $\beta$  III-tubulin isotype III clone SDL.3D10, Sigma, Cat# T8660
  - Rabbit anti-GFAP, Dako, Cat# M0761

**Protocol**

**Notes:**

- *We do not recommend re-freezing supplements and medium provided with the Neurons Starter Kit.*
- *We do not recommend cryopreserving Neurons Starter Kit cultured neurons.*

**Simplified diagram showing key steps in the process of culturing Applied StemCell neurons**



**1. Handling Upon Receiving**

Neurons Starter Kit is shipped on dry ice. The components are packed in zip-lock bags. A vial containing cryopreserved neuronal precursors is packed in a small transparent bag which is buried in dry ice. Upon receiving the product, check integration of the package and the presence of dry ice (contact Applied StemCell if the integrity of the package has been compromised, e.g. no dry ice in the package).

Cells can be plated immediately after arrival or transferred into liquid nitrogen. Do not remove the vials from dry ice during transportation to storage units. Immediately transfer components of the Neurons Starter Kit (especially the cryopreserved neuronal precursors) to storage units, avoiding exposure to room temperature.

After arrival, properly store the kit's components as follows:

Component	Storage
Neuronal precursors	Liquid Nitrogen
Neuron Maturation Medium	-20°C
Supplement A	-20°C

## 2. Procedure

This procedure has been extensively tested with Applied StemCell neurons and Applied StemCell Neuron Maturation Medium. The user should strictly follow provided protocol. The user assumes all responsibility for the failure of the experiment should there be any deviation from this procedure. Please read and understand the entire protocol before proceeding with the maturation procedure.

### 2.1 Coating Cell Culture Vessels with Poly-L-ornithine and Laminin

Cell culture vessels should be coated one day before or on the day of plating the cells. Please read producer's manual for handling of Poly-L-ornithine and Laminin.

- 2.1.1. Prepare stock solution of poly-L-ornithine (10 mg/mL) by dissolving the powder in sterile cell culture grade water. The stock solution should be stored at -20°C.
- 2.1.2. Thaw stock solution of laminin (1 mg/mL, Life Technologies) on ice.
- 2.1.3. Prepare working solution of poly-L-ornithine in sterile cell culture grade water at a final concentration of 20 µg/mL.
- 2.1.4. Add poly-L-ornithine solution into desired cell culture vessel to cover the vessel's bottom entirely (see Table 1).
- 2.1.5. Distribute the solution evenly and incubate vessels in the cell culture incubator for 2 hr (37°C/ 5% CO<sub>2</sub>/ humidity control).
- 2.1.6. Rinse vessels two times with cell culture grade water. Pipette water gently toward the corner of the vessel to avoid mechanical removal of poly-L-ornithine coating.
- 2.1.7. Prepare working solution of laminin in sterile cell culture grade water at a final concentration of 10 µg/mL.
- 2.1.8. Aspirate water from the vessels and add laminin solution to cover well the vessel's bottom. Incubate in the cell culture incubator (37°C/ 5%CO<sub>2</sub>/ humidity control) for 2 hours.
- 2.1.9. If not used immediately, store coated vessels at 4°C in laminin solution (up to 4 days).
- 2.1.10. Pre-warm vessels at 37°C before use.
- 2.1.11. Aspirate laminin just before seeding DOPA precursors. Do not let the surface dry.

**Table 1. Recommended volumes of coating reagents in various vessels.**

Vessel Type	P-L-ornithine	Laminin
96 well plate	50 µL/well	50 µL/well
4 or 24 well plate	250 µL/well	250 µL/well
35 mm dish	1.5 mL	1.5 mL
60 mm dish	2.5 mL	2.5 mL

## 2.2 Thawing and Culturing Cryopreserved Neurons

All steps described here must be performed in accordance with aseptic cell culture standards. All media and vessels used for cell culture must be pre-warmed to 37°C prior to use. For the entire process of neuron maturation, one type of complete medium is required: Neuron Maturation Medium + Supplement A.

2.2.1 One day before thawing precursor cells, place the 50 mL Neuron Maturation Medium bottle in 2°- 8°C fridge overnight.

*Note: Once thawed, Neuron Maturation Medium can be stored at 2°- 8°C for up to 3 weeks.*

2.2.2 On the day of thawing cryopreserved neurons, transfer a 25 mL aliquot of Neuron Maturation Medium into a 50 mL conical tube and add 25 µL of Supplement A from vial 1 (pre-thawed on ice) to make the complete medium.

2.2.3 Transfer a 5 mL aliquot of the complete medium prepared in step 2 into a 15 mL conical tube and pre-warm at 37°C. This aliquot will be used for recovery of precursor cells from frozen stock.

2.2.4 Prepare another aliquot of the complete medium in accordance with volumes required for cell culture vessels utilized (see Table 2). Only take enough medium to be utilized for cell culture that day and pre-warm at 37°C. Place the rest of the complete medium back to 2°- 8°C fridge.

**Table 2. Recommended volumes of medium in various vessels**

Vessel Type	Medium Volume
96 well plate	100 µL/well
4 or 24 well plate	500 µL/well
35 mm dish	2 mL
60 mm dish	5 mL

2.2.5 Shortly before thawing the cells, place pre-warmed medium and vessels in a biosafety cabinet.

2.2.6 To thaw cryopreserved neurons, remove one vial from storage unit and place immediately onto dry ice (the vial must be buried in the dry ice).

2.2.7 Bring dry ice container with the vial to a 37°C water bath and immerse the vial in the bath (up to 2/3 of the vial) and thaw cells until only a small piece of ice is still visible (approximately one minute).

*Note: Do not shake the vial during thawing.*

2.2.8 Immediately bring the vial to the biological cabinet, spraying it thoroughly with 70% ethanol and wiping with an autoclaved paper towel.

2.2.9 Remove cells from the vial using a p1000 micropipette (or serological pipette) and transfer drop-wise while swirling into 15 mL conical tube containing 5 mL of pre-warmed complete medium (step 3). Wash the vial with 1 mL medium from the 15 mL conical tube and transfer it back to the tube.

*Note: Do not mix cells up and down and avoid generation of bubbles.*

- 2.2.10 Centrifuge cells at 400 x g for 5 minutes at room temperature.
- 2.2.11 Aspirate the medium very carefully using a vacuum (or pipette if preferred), leaving only a drop of liquid in the tube. Take extra care not to remove or disturb the cell pellet during aspiration of medium.
- 2.2.12 Using a p1000 micropipette, add 1 mL of the complete medium (step 4) into the tube and gently re-suspend cells by pipetting up and down 4-6 times.
- 2.2.13 Remove a 10  $\mu$ L aliquot of cell suspension and mix it with 10  $\mu$ L of Trypan blue solution.
- 2.2.14 Count the cells.
- 2.2.15 Take the appropriate volume of pre-warmed complete Neuron Maturation Medium and add to the tube with the cells.
- 2.2.16 Aspirate laminin solution from pre-warmed cell culture vessel and seed neurons at a density ranging from  $4 \times 10^4$  to  $8 \times 10^4$  live cells/cm<sup>2</sup>. See Table 3 for your convenience.

**Table 3. Recommended seeding densities for Applied StemCell neurons in various types of cell culture vessels. Range: from low to high.**

Vessel	Surface/Well	Medium Volume	Density (Cells)
96-well plate	0.33 cm <sup>2</sup>	100 $\mu$ L/well	$1.3 \times 10^4 - 2.6 \times 10^4$
4-well plate	2 cm <sup>2</sup>	500 $\mu$ L/well	$8 \times 10^4 - 1.6 \times 10^5$
35 mm dish	10 cm <sup>2</sup>	2 mL	$4 \times 10^5 - 8 \times 10^5$
60 mm dish	20 cm <sup>2</sup>	5 mL	$8 \times 10^5 - 1.6 \times 10^6$

- 2.2.17 Distribute cells evenly and place vessels in the cell culture incubator (37°C/ 5% CO<sub>2</sub>/humidity control). This is **Day 0**; Medium should be changed ever other day.
- 2.2.18 Monitor the cells' survival and attachment the following day (**Day 1**).
- 2.2.19 Change complete Neuron Maturation Medium on **Day 2**. Medium change should be done slowly (drop wise), pointing the pipette tip toward the wall of cell culture vessel.
- 2.2.20 Change medium every other day. Continue maturation of neurons for 8 days.

*Note: Cells can be differentiated for up to 5 weeks. However, prolonged culture will have increased population of GFAP positive astrocytes.*